



# FRESH BAIT



Individually packed, 10-gram ready-to-use paste formulation for rat (*Rattus norvegicus*) and mouse (*Mus musculus*) control indoor and outdoor. The soft, oily paste with its intensive smell and excellent taste attracts rodents even in those places where they can easily find other food as well.

**ACTIVE INGREDIENT:** 0.0027 % bromadiolone

### PHYSICAL PROPERTIES, APPEARANCE:

**APPEARANCE:** red coloured paste  
**SIZE:** ~ 55 x 70 mm  
**ODOUR:** characteristic to paraffin  
**SOLUBILITY:** insoluble in water  
**FLAMMABILITY:** combustible

### TOXICOLOGY

ACUTE ORAL (rat) LD<sub>50</sub> > 2 000 mg/kg b.w.  
ACUTE DERMAL (rat) LD<sub>50</sub> > 2 000 mg/kg b.w.

**Antidote:** Vitamin K1

### ADVANTAGES

- Contains second generation anticoagulant active ingredient.
- A single consumption is enough for the elimination of rodents.
- Low active ingredient content.
- The fresh taste makes it very attractive to rodents.
- Coloured bait with bitter agent to avoid accidental consumption by humans and domestic animals.

### PRODUCT FOR PROFESSIONAL USE

**STABILITY/ SHELF LIFE:**  
2 years from date of manufacture, if stored in original unopened packaging in a dry and cool place

### PACKING:

KS-11156	 5 kg	 64 buckets
KS-11157	 25 kg	 24 cartons





# FRESH BAIT



## INSTRUCTIONS FOR USE:

Place baits manually in the rodent infested area. The method of deployment is dependent on the particular circumstances, however priority should be given to exclude non-target exposure as much as possible.

For rat infestations use bait points up to 120 g bait in bait stations. In case of baiting boxes, use the built in hangers/spikes for fixing the sachets. Ideally bait boxes should be fixed to the ground.

Against rats place bait points 10 metres apart for low infestations and 5 metres for high infestations.

For mouse infestations use bait points of up to 40 g bait in bait stations (mouse boxes) or covered bait points.

Bait may also be placed on trays under a tile or located where access by non-target organisms is restricted. In this case – as far as possible – fix the bait sachets (e.g. with wire), so that they cannot be dragged away (or dispersed) by rodents.

Against mice place bait points 10 metres apart for low infestations and 5 metres for high infestations.

The product must never be placed indiscriminately.

Death of the rodents starts after 3 – 5 days from the beginning of consumption. Make regular inspections of the bait points (every 7 - 10-days) and replace any bait that has been eaten by rodents, damaged by water or contaminated by dirt. If bait is totally consumed then each bait point should be replenished.

## Warning:

Read the instructions before use!

It can only be used for rodent control and according to the instructions for use!

The placed out rodenticide bait must always be labelled with a warning sign. After placing the baits wash your hands carefully with warm water and soap.

Put the bait in areas where children, domestic animals, livestock and birds do not have access to them. The preparation is toxic for mammals and birds. Dogs, cats, pigs and other predator and/or scavenger animals may get poisoned if they eat rats or mice that died or had become ill from the rodenticide. It is forbidden to use the preparation to control non-target animals. It is forbidden to let the preparation enter surface waters.

The safety data sheet is available for professional pest controllers.

