



# GRAIN BAIT



Ready-to-use, specially coated grain bait formulation with second generation anticoagulant active ingredient content for rats and mice. Available in different bulk packages for use in bait boxes or in aroma permeable sachets to place without opening.

**ACTIVE INGREDIENT:** 0.0027 % bromadiolone

### PHYSICAL PROPERTIES, APPEARANCE:

**APPEARANCE:** coarse grain  
**COLOUR:** pink dyed  
**ODOUR:** no odour  
**SOLUBILITY:** insoluble in water  
**FLAMMABILITY:** combustible

### TOXICOLOGY

ACUTE ORAL (rat) LD<sub>50</sub> > 2 000 mg/kg b.w.  
 ACUTE DERMAL (rat) LD<sub>50</sub> > 2 000 mg/kg b.w.

**Antidote:** Vitamin K1

### ADVANTAGES

- Contains second generation anticoagulant active ingredient.
- A single consumption is enough for the elimination of rodents.
- The scientifically tested aroma permeable packaging lets rodents feel the scent of the bait, thus attracting rats and mice as effectively as if it was opened.
- The sachet allows for precise dosing and makes it easy to mark, control and measure consumption, resulting in maximum cost-effectiveness.
- Coloured bait with bitter agent to avoid accidental consumption by humans and domestic animals.

### PRODUCT FOR PROFESSIONAL USE

**STABILITY/ SHELF LIFE:**  
2 years from date of manufacture, if stored in original unopened packaging in a dry and cool place

### PACKING:

KS-11153	 4 kg	 80 buckets
KS-11154	 400x25 g sachet	 32 cartons





# GRAIN BAIT



## INSTRUCTIONS FOR USE:

### GRAIN BAIT IN SACHET

Place the sachet containing the rodenticide bait - without opening the filter-paper covering- on a suitable tray or in baiting box to the locations visited by mice and rats, near the runs and assumed hiding places of rodents. Fix the sachet inside the container (spike, locking rods).

Amount to be used depending on the degree of infestation:

- 1- 4 pcs filter-paper covered sachets per 5 m<sup>2</sup> to control mice,
- 8- 10 pcs filter-paper covered sachets for every 7-10 metres to control rats.

### GRAIN BAIT IN BULK

Place the rodenticide grain bait in the container intended for this purpose to the locations visited by mice and rats, near the runs and assumed hiding places of rodents. Open the sachet before use. In the case of bulk packs (not packed in sachet) use a suitable respirator when pouring the product.

Amount to be used depending on the degree of infestation:

- 50-100 grams per 5 m<sup>2</sup> to control mice,
- 200-250 grams for every 7-10 metres to control rats.

Rodents start to die 4-5 days after consumption. Check the bait placed out every 4-7 days and replace the consumed and/or carried away bait. Remove contaminated or expired baits and replace them by fresh sachets. Continue placing out the baits until rodents disappear completely. If the sachets remain intact but the rodents are still present, transfer the rodenticide to somewhere else. Depending on the rate of their infestation and the size of the area you can expect extermination of all the rodents in 2-3 weeks.

### Warning:

Read the instructions before use!

It can only be used for rodent control and according to the instructions for use!

The placed out rodenticide bait must always be labelled with a warning sign. After placing the baits wash your hands carefully with warm water and soap.

Put the bait in areas where children, domestic animals, livestock and birds do not have access to them. The preparation is toxic for mammals and birds. Dogs, cats, pigs and other predator and/or scavenger animals may get poisoned if they eat rats or mice that died or had become ill from the rodenticide. It is forbidden to use the preparation to control non-target animals. It is forbidden to let the preparation enter surface waters.

The safety data sheet is available for professional pest controllers.

